


Patient Guideline for Micro Leuprolide or Leuprolide Trigger

1. Refrigerate vial. Store in center of refrigerator, avoid storing in the door or back.
2. Check that solution is not discolored.
3. Discard medication after expiration on vial. Overfill is given to provide extra, therefore it is OK if there is liquid remaining in vial.

Directions for injecting:

1. Wash your hands.
2. Take leuprolide vial out of refrigerator. Let it come to room temperature for about 10-15 minutes. This will allow less burning/stinging.
3. Set out alcohol swabs, syringe, and Sharps container.
4. Cleanse injection site with alcohol swab. Abdomen is preferred, but front of thigh and the back/side of the upper arm are other possible spots.



5. Let the alcohol air dry completely before injecting. This will also allow less burning/stinging.
6. Remove the sterile sticker from the vial and wipe rubber top of vial with alcohol swab.
7. Take syringe and pull back the air equal to the dose amount you will be injecting.
8. At a 45 degree angle, puncture rubber stopper with syringe and inject air.
9. Keeping the eye of the needle in the liquid, draw up "units" or "ml" needed for dose.
10. Flick out any air bubbles.
11. Pinch about a 2 inch area of the skin where injecting (always rotate injection sites). Inject at a 45 to 90 degree angle and insert entire length of the needle, release the skin, and depress the plunger slowly until all medication is injected. Remove the syringe quickly and gently.
12. Dispose of syringe in Sharps container.